



FLUENZ TETRA NASAL SPRAY FLU VACCINE

FOR 2- AND 3-YEAR-OLDS AND CHILDREN AT RISK

Aged 4 to 17 years –should be given at school. (All children in Reception class and in school years 1, 2, 3 and 4 and Children up to 17 who are 'at risk' from flu, such as children with long-term health conditions)

Any child 2-9 in a Clinical Risk Group, who has not received the Seasonal Influenza Vaccine before, should have a 2nd dose of Fluenz 4 weeks after the 1st dose.

Name of the child:

Date of Birth:

Please let the nurse know if any of these contra indications apply to your child: Is your child:

1. Less than 2 years of age
2. Severely immunodeficient due to any condition e.g. leukaemia, lymphoma, high dose steroids, and HIV infection.
3. On any immunosuppressive therapy
4. On any salicylate therapy in medicines for pain relief
5. Allergic to eggs, gentamicin, or gelatine
6. Wheezing today or severe blocked nose
7. Severe Asthma, sign step 4 or above
8. Had another live vaccine e.g. MMR, Yellow Fever, BCG within the last 4 week
9. Pregnant or likely to be in the next month, or breast feeding.
10. In the next 2 weeks is likely to encounter anyone who has a lowered immunity.

Your child may experience nasal congestion, decreased appetite, headache, malaise, temperature.

Very rarely they may get itchy skin, rash, muscle ache, nose bleeds or more severe reaction. SO PLEASE STAY ON THE PREMISES FOR 10 MINUTES AFTER THE VACCINATION.

Signed:

Date:

Office use only: Vaccine Given:

Prescription Done: